

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US05/001807

International filing date: 20 January 2005 (20.01.2005)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/537,864
Filing date: 21 January 2004 (21.01.2004)

Date of receipt at the International Bureau: 03 March 2005 (03.03.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

1286783

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

February 17, 2005

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/537,864

FILING DATE: *January 21, 2004*

RELATED PCT APPLICATION NUMBER: *PCT/US05/01807*



Certified by

Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office

TITLE: **CHEMICAL INHIBITORS OF SOLUBLE
ADENYLYL CYCLASE (sAC)**

INVENTORS: **Jochen Buck and Lonny R. Levin**

DOCKET NO.: **19603/4650 (CRF D-3373-01)**

PROVISIONAL PATENT APPLICATION

**Bicarbonate responsive 'soluble' adenylyl cyclase defines a nuclear
cAMP microdomain**

Introduction

Cyclic AMP is a nearly ubiquitous second messenger molecule that affects a multitude of cellular functions. In mammalian cells, two classes of adenylyl cyclase generate cAMP. Transmembrane adenylyl cyclases (tmACs) are tethered to the plasma membrane and regulated by heterotrimeric G proteins in response to hormonal stimuli [reviewed in (Hanoune and Defer, 2001)]. A second source of cAMP, the more recently described 'soluble' adenylyl cyclase (sAC), resides in discrete compartments throughout the cell (Zippin et al., 2003) and is regulated by the intracellular signaling molecules bicarbonate (Chen et al., 2000) and calcium (Jaiswal and Conti, 2003; Litvin et al., 2003).

Cyclic AMP elicits its cellular effects by activation of three known classes of effector proteins; Exchange Proteins Activated by cAMP (EPAC), Cyclic Nucleotide Gated Ion Channels (cNGC), and Protein Kinase A (PKA). A subset of these targets reside at the plasma membrane, where they exist in macromolecular signaling complexes which also include a receptor [a G protein coupled receptor (GPCR)], its transducing G-protein, and the source of cAMP, a tmAC isoform (Davare et al., 2001). The cAMP generated by tmACs acts locally (Rich et al., 2000; Rich et al., 2001; Zaccolo and Pozzan, 2002), most likely restricted by phosphodiesterase (PDE) 'firewalls' (Zaccolo and Pozzan, 2002) which define the limits of these cAMP signaling microdomains. But targets of cAMP do not solely reside at the plasma membrane. EPAC is localized to the nuclear membrane and mitochondria (Qiao et al., 2002), and PKA is tethered throughout the cell by a class of proteins called A-kinase anchoring proteins (AKAP) (Michel and Scott, 2002). The observation that cAMP does not diffuse

far from tmACs (Bacskai et al., 1993; Zaccolo and Pozzan, 2002) reveals there must be another source of cAMP modulating the activity of these distally localized targets.

‘Soluble’ adenylyl cyclase (Buck et al., 1999) is widely expressed in mammalian cells (Sinclair et al., 2000). Unlike tmACs, sAC is G protein insensitive (Buck et al., 1999), and among mammalian cyclases, it is uniquely responsive to intracellular levels of bicarbonate (Chen et al., 2000). The ubiquitous presence of carbonic anhydrases ensures that the intracellular bicarbonate concentration (and sAC activity) will reflect changes in pH (Pastor-Soler et al., 2003) and/or CO₂. Because CO₂ is the end product of energy producing metabolic processes, sAC is poised to function as a cell’s intrinsic sensor of metabolic activity (Zippin et al., 2001). sAC possesses no transmembrane spanning domains (Buck et al., 1999) and is distributed to subcellular compartments containing cAMP targets (Zippin et al., 2003) which are distant from the plasma membrane. sAC was also found localized inside the mammalian cell nucleus (Zippin et al., 2003).

To evaluate how sAC-generated cAMP might differ from the second messenger generated by tmAC’s, we explored a prototypical cAMP dependent pathway, PKA dependent phosphorylation of cAMP response element (CRE) Binding protein (CREB) (De Cesare and Sassone-Corsi, 2000). In a widely accepted signal transduction paradigm, extracellular signals (i.e., hormones and neurotransmitters) affect CREB family phosphorylation by stimulation of plasma membrane bound tmACs. The generated cAMP activates nearby PKA, and the liberated catalytic subunit then appears to translocate through the cytoplasm to phosphorylate and activate CREB proteins residing inside the nucleus (Hagiwara et al., 1993; Riabowol et al., 1988b). Intracellular signals, such as metabolic activity, also modulate CREB phosphorylation in a cAMP-

dependent manner (Daniel et al., 1998; Singh et al., 2001; Trumper et al., 2002), but the mechanism has yet to be established. Localization of sAC inside the nucleus, in close proximity to the CREB family proteins, and its regulation by calcium and bicarbonate, suggested that sAC might be responsible for modulating CREB activity in response to intracellular signals.

In this report, we demonstrate the existence of a nuclear cAMP signaling microdomain that mediates bicarbonate dependent activation of the transcription factor CREB. Bicarbonate activation of CREB represents an example of a mammalian cAMP dependent pathway solely modulated by intrinsic cellular signals. This nuclear cAMP signaling cascade functions independently from the classically defined mechanisms leading to CREB activation, demonstrating that cAMP is a locally acting second messenger which can work autonomously in different compartments within a single cell.

Results

Bicarbonate induces CREB phosphorylation

Bicarbonate treatment of cells uniquely activates sAC (Chen et al., 2000) while activation by G_s proteins or forskolin only stimulates tmACs; therefore, these agents can be used to differentially stimulate the two classes of mammalian adenylyl cyclase. To determine whether sAC activation would elicit PKA activation of CREB, a well-characterized target of tmAC generated cAMP, we treated cells with bicarbonate and measured PKA dependent phosphorylation of CREB using antisera specific for the PKA (Ser133) phosphorylated form of CREB. Hormonal stimulation of CREB transcription factors, acting through tmACs, reaches its peak in 30 minutes (Hagiwara et al., 1992). Treatment of COS7 cells with forskolin, which will activate the total cellular pool of tmACs, stimulated nuclear immunofluorescent staining and Western blot immunoreactivity using the phospho-specific antisera (Figures 1A,B). Treatment of COS7 cells for the same amount of time (30 minutes) with bicarbonate also resulted in CREB phosphorylation (Figures 1A,C). These increases in phospho-CREB immunostaining were inhibited by pretreatment with H89, confirming the involvement of PKA (Figures 1A-C). Overexpression of sAC led to an increase in basal CREB phosphorylation (Figure 1B+1C, lane 4) suggesting that sAC-generated cAMP was sufficient to activate CREB. Consistent with its bicarbonate responsiveness (Chen et al., 2000), sAC overexpressing COS7 cells displayed enhanced bicarbonate-dependent CREB phosphorylation (Figure 1C, lanes 2 and 5) which was also blocked by H89 (Figure 1C). The ability of either bicarbonate or forskolin to induce CREB phosphorylation reveals that CREB represents a downstream target of both tmAC and sAC generated cAMP.

Time course of bicarbonate induced CREB phosphorylation

We directly compared the time course of CREB activation in response to a hormonal activator of tmACs, PGE₂, versus the specific sAC activator, bicarbonate, in a liver cell line (Hagiwara et al., 1992). Phosphorylation of CRE binding proteins in response to bicarbonate occurred rapidly; increases in phospho-CREB were detected within 2 minutes, the earliest time tested (Figure 2A). In contrast, PGE₂ (Figure 2B) or forskolin (data not shown) stimulation of CREB phosphorylation was detectable only after 5 minutes, consistent with published reports (Hagiwara et al., 1992). The longer activation kinetics following PGE₂ or forskolin stimulation is thought to reflect the time required for translocation of PKA catalytic subunit into the nucleus from the plasma membrane where it was activated by a hormonally modulated tmAC (Hagiwara et al., 1993). In addition to being more rapid, the peak intensity of phosphorylation was higher with bicarbonate treatment. The different kinetics and intensity of CREB activation by bicarbonate and PGE₂ reveal that while sAC and tmACs may affect overlapping substrates, they may participate in distinct signal transduction cascades.

CREB, sAC, and PKA co-exist in the nucleus

Because CREB family members and sAC (Zippin et al., 2003) reside inside the nucleus, we reasoned the accelerated kinetics and intensity of bicarbonate-induced CREB activation could occur if sAC and CREB co-existed in a signal transducing complex. A complete nuclear cAMP signaling cascade capable of phosphorylating CREB family proteins requires the presence of the cAMP-responsive PKA holoenzyme. Both catalytic and regulatory subunits of PKA have been immunologically (Jungmann

et al., 1988; Kuettel et al., 1985; Yang et al., 1998) and biochemically (Byus and Fletcher, 1982; Constantinescu et al., 1999; Murray et al., 1985; Zhang et al., 1996) detected inside the nucleus, and nuclear localization of PKA holoenzyme has been described in lower eukaryotes (Griffioen et al., 2000), but the nuclear presence of PKA regulatory subunit, and therefore cAMP-responsive holoenzyme, has been questioned. We repeated and extended the immunological examination of regulatory subunit localization and confirmed that PKA resides inside the nucleus of the human liver cell line, Huh7 (Figure 3A), suspension HeLa cells (Figure 3B-C), and in a subset of cells within sectioned liver tissue (Figure 4A). Confocal microscopy of Huh7 and HeLa cells using polyclonal and monoclonal antibodies, recognizing PKA regulatory subunit isoforms (RI α , RII α), revealed distinctive cytoplasmic staining in accordance with accepted dogma (Alto et al., 2002) (Figure 3A-C), but these regulatory subunit isoforms were also detected inside the nucleus (Figure 3A-C and 4A). In the case of suspension HeLa cells, it should be stressed that these optical slices were selected to illustrate the intranuclear staining of PKA. Slide preparation and imaging constraints cause PKA cytoplasmic staining to appear as a thin layer surrounding the nucleus (Figure 3B-C, arrows marked A) and within the expanse of cytoplasm stretching out as these suspension cells adhere to the coverslip (Figure 3B-C, arrows marked B).

Nuclear staining of each isoform was distinct. RII α was present in a diffuse pattern throughout the nucleus with small areas of enrichment (Figure 3A, middle panels and 3B), whereas RI α was distributed in the nucleoplasm but more enriched in nucleoli (Figure 3A, right panels and 3C). RII α was also detected in the nuclei of a subset of rat liver primary hepatocytes (Figure 4A). Consistent with our previously published data (Zippin et al., 2003), sAC was also present in the nuclei of Huh7 cells

(Figure 3D) and a subset of rat liver hepatocytes (Figure 4). PKA, sAC, and phosphorylated CREB seem to be co-ordinately localized; the subset of nuclei in rat liver hepatocytes and Huh7 cells (data not shown) positive for sAC protein (Figure 4A and B, arrow A) also contained R subunit (Figure 4A, arrow A) and CREB phosphorylation (Figure 4B, arrow A), whereas nuclei not enriched for sAC displayed neither R subunit nor CREB phosphorylation (Figure 4A and B, arrow B). Rat liver hepatocytes positive for sAC, PKA, and phospho-CREB represent approximately 10% of total hepatocytes, and we have not yet identified any consistency with known liver anatomy. These data demonstrate that nuclei contain all the components of a cAMP signalling cascade and suggest that sAC generated cAMP is positioned to activate nuclear PKA holoenzymes to phosphorylate CREB proteins.

Isolated nuclei contain components of a cAMP signaling microdomain

Bicarbonate treatment of whole cells leads to rapid induction of CREB phosphorylation (Fig. 2). To test whether the nuclear localized sAC and PKA were responsible for this bicarbonate induced CREB activation, we prepared isolated nuclei from suspension HeLa cells, a cell line with well established protocols for the isolation and enrichment of nuclei. Cells were lysed using digitonin and nuclear preparations were purified by density centrifugation through an Optiprep gradient. Western analyses of the same cell equivalents from each fraction using cellular markers for different subcellular compartments (histone H3, Na/K ATPase α 1 subunit, cytochrome c oxidase subunit III (COX), and β -tubulin) confirmed that the nuclear fractions (P2) were positive for nuclear markers (histone) with undetectable levels of plasma membrane (Na/K ATPase), mitochondrial (COX), or cytoplasmic (tubulin) contamination (Figure

5A). To confirm that the P2 fraction did not contain any detectable mitochondria, a possible source of both sAC and PKA contamination, we overloaded the P2 fraction, but COX antigen was still not detected (data not shown). Visual inspection and DAPI fluorescence confirmed that the final preparation was enriched for intact nuclei (Figures 5B and 5C), and as expected, isolated nuclei contained both CREB and sAC proteins by immunocytochemistry (Figure 5B) and by Western blotting (Figure 5D)

Consistent with staining patterns shown above (Figures 3A-C), RII α immunostaining was present throughout the nucleus, while RI α appeared enriched within the nucleolus (Figure 5C). PKA RI α and RII α were also detected by Western analysis as a single band of the predicted molecular weight in the P2 lysate, using monoclonal and polyclonal antibodies, (Figure 5E) confirming the specificities of these antibodies for immunostaining. Because the staining patterns of isolated HeLa cell nuclei (Fig. 5C) reflected the immunostaining pattern observed in intact HeLa cells (Fig. 5B-C), we concluded that the isolation and enrichment of nuclei had little effect on nucleoplasm architecture.

sAC represents the only source of cAMP detectable in isolated nuclei

We previously demonstrated that sAC activity was present in COS7 cell nuclei (Zippin et al., 2003). We now show that bicarbonate responsive sAC is the only source of cAMP in nuclei isolated from suspension HeLa cells. Whereas forskolin potently stimulates cAMP production in whole cell lysates (Figure 6A), there was no significant increase in cAMP elicited by forskolin in isolated nuclei (Figure 6B). There was a significant level of basal adenylyl cyclase activity in isolated nuclei, which was stimulated by bicarbonate addition (Figure 6B). Both the bicarbonate stimulated and

basal activities were inhibited by sAC-selective inhibitors (Figure 6C,D). We have identified several sAC inhibitors (Figure 6C), inert towards tmACs (Figure 6D), in a screen of a combinatorial chemical library (K. H., T.N. Litvin, E. Hyde, J.H.Z., L.R.L., and J.B., in preparation). In the presence of two representative, structurally unrelated inhibitors (KH1 and KH2 each display an IC_{50} of $\leq 10 \mu M$ towards recombinant human sAC protein), the cAMP generated in the presence of bicarbonate in P2 nuclei was reduced to a level below that of basal. These results indicate that in addition to mediating the bicarbonate-induced increase in cAMP in isolated nuclei, sAC is also responsible for the observed basal adenylyl cyclase activity.

Bicarbonate induces CREB phosphorylation in isolated nuclei

CREB phosphorylation in isolated nuclei was assayed by immunocytochemistry using phospho-CREB specific antisera (Figure 7A-B). Nuclei incubated in the presence of either bicarbonate or cAMP displayed at least a two-fold rise in the percentage of phospho-CREB positive nuclei relative to untreated nuclei (basal) (Figure 7B). As expected, due to the lack of tmACs in isolated nuclei, the number of nuclei positive for CREB phosphorylation was unaffected by forskolin. These data demonstrate that a bicarbonate-responsive signaling cascade leading to CREB phosphorylation is wholly contained within the mammalian cell nucleus. In contrast, the hormone and forskolin-responsive tmAC defined cascade is only functional in a whole cell context.

Nuclear sAC activates CREB via nuclear PKA

To facilitate the use of pharmacological reagents to further evaluate bicarbonate-induced CREB phosphorylation, we monitored CREB phosphorylation by Western analysis (Figure 7C-D). Similar to our observations using immunocytochemistry, treatment of isolated nuclei with bicarbonate or 8-Br-cAMP elicited a 27-fold or 30-fold increase in CREB phosphorylation, respectively (Figure 7C). Once again, forskolin, which had a potent effect in a whole cell context (Figure 1B), elicited no significant stimulation of CREB phosphorylation in isolated nuclei (Figure 7D).

We next confirmed that the effects of bicarbonate on CREB phosphorylation were mediated by nuclear sAC and PKA. CREB phosphorylation induced by bicarbonate was substantially reduced by the PKA inhibitors, H89 (50%) and RpcAMPs (70%) (Figure 7C), revealing the involvement of cAMP-responsive PKA holoenzyme. The chemical inhibitors effective at blocking sAC generated cAMP accumulation (KH1 and KH2, Figure 6B) were also effective at preventing bicarbonate induced CREB phosphorylation (Figure 7D) demonstrating, once again, that sAC is responsible for the bicarbonate-stimulated cAMP-dependent, phosphorylation of CREB in the mammalian cell nucleus.

Discussion

Most cellular pathways in eukaryotic cells are impacted by cAMP. Effectors of cAMP mediate processes at both the plasma membrane and multiple, distinct intracellular sites. It has been widely assumed that cAMP is generated exclusively at the plasma membrane by G protein-regulated tmACs, and the second messenger then diffuses from the cell membrane through the cytosol to its intracellular targets. However, FRET-based (Bacskai et al., 1993; Zaccolo and Pozzan, 2002) and biochemical (Rich et al., 2000; Rich et al., 2001) methods for observing intracellular cAMP concentrations reveal that the second messenger generated by tmACs does not diffuse far from its site of synthesis. We have recently demonstrated that sAC is localized at multiple, subcellular compartments throughout the cell, including mitochondria, centrioles, the mitotic spindle, the mid-body and the nucleus (Zippin et al., 2003), each of which contain targets of cAMP. These data suggest that the cell may contain multiple, independently modulated cAMP signaling microdomains; targets near the plasma membrane would depend upon tmACs for second messenger generation, while targets inside the cell would be modulated by sAC generated cAMP (Wuttke et al., 2001; Zippin et al., 2003). We now provide data supporting this hypothesis by demonstrating the existence of a sAC-defined nuclear cAMP signalling microdomain which can lead to CREB activation.

The nuclear cAMP signaling cascade induced by bicarbonate produced a rapid activation of CREB family members in both whole cells and nuclei, while PGE₂ and forskolin, tmAC specific activators, produced a delayed response exclusively in whole cells. Therefore, cAMP-mediated activation of CREB family members by tmACs and sAC proceed via independent pathways. CREB activation by hormones or

neurotransmitters via tmACs apparently requires time for movement of PKA catalytic subunit from the plasma membrane into the nucleus (Hagiwara et al., 1993; Riabowol et al., 1988a). This delayed activation is consistent with hormonal control of gene expression providing a long-term response to predominantly sustained extracellular signals (Bailey et al., 1996). In contrast, the newly described nuclear sAC activation pathway proceeds rapidly without requiring the translocation of any constituent. In this regard, the sAC nuclear microdomain is capable of responding quickly to subtle fluctuations in intrinsic signals, such as local intracellular concentrations of bicarbonate and calcium.

In tissues, sAC is not present within the nucleus of every cell. In liver, sAC appears to be predominantly extranuclear but enriched in a subset of the nuclei (Figure 4, arrows A). PKA holoenzyme appears to be enriched within the same subset of nuclei (Figure 4A, arrow A), and interestingly, these are the nuclei which are also positive for CREB phosphorylation (Figure 4B, arrow A). The presence of both positive and negative nuclei for sAC, PKA, and CREB phosphorylation in the same tissue suggests that there may be coordinated regulation of the presence of this newly described nuclear signaling microdomain.

The demonstration that bicarbonate treatment of whole cells leads to activation of the CREB family of transcription factors reveals that bicarbonate itself induces a signal transduction cascade. Cellular bicarbonate levels reflect intracellular pH as well as CO₂ generation (Bevensee et al., 2000); therefore, bicarbonate signaling pathways would respond to a wide variety of cellular transitions. Immunostaining revealed that sAC is present at mitochondria, centrioles, mitotic spindles, and mid-bodies (Zippin et al., 2003), suggesting the existence of multiple cAMP signaling microdomains within a

single cell. A remaining challenge will be to determine whether sAC molecules in these different microdomains are subject to independent and unique modes of regulation, permitting a variety of distinct responses independently mediated by the same second messenger.

Materials and Methods

Cell Growth and Transfections

All cell lines were grown in DMEM (44 mM sodium bicarbonate) supplemented with 10% FBS. Where indicated, cells were transfected using Lipofectamine 2000 (Invitrogen) per manufacturer's instructions. Cells were incubated with DNA for 5 hours in OPTI-MEM and then switched to normal media. Bicarbonate starvation was conducted by changing media to bicarbonate-free DMEM (44 mM Hepes) supplemented with 10% FBS for at least 1 hour at 37°C under ambient air conditions. Bicarbonate stimulation consisted of returning cells to normal bicarbonate containing media and placing them in a 5% CO₂ incubator. For PGE₂ or forskolin stimulation, cells were grown in normal media under 5% CO₂. Stimulation was accomplished by replacing media with normal media containing 1 μM PGE₂ or 10μM forskolin. For Western analysis, cells were lysed immediately by direct addition of SDS sample buffer.

Immunocytochemistry

Cells or nuclei were washed in PBS and fixed for either 30 minutes in 4% PFA and permeabilized in 0.1% Triton X-100 or fixed for 15 minutes in 2% PFA and permeabilized in 0.05% Triton X-100. Liver from adult rat was rapidly excised, placed between two thinly sliced pieces of bovine liver, and snap frozen in isopentane cooled with liquid nitrogen. Cryosections (6 μM thick) were collected on superfrost slides (Fisher Scientific) and stained within 1 day of sectioning. Tissue was fixed for 30 minutes in 4% PFA and permeabilized in 0.1% Triton X-100 for 15 minutes. All samples were blocked in 2% BSA for at least one hour. Cells or tissues were stained with anti-sAC R41 or R52 biotinylated monoclonal antibodies or R21 monoclonal

antibody (1:100) generated against human sAC₁ antigen as previously described (Zippin et al., 2003), anti-PKA regulatory subunit (RI α and RII α) polyclonal antisera (1:100, Chemicon International and Cedarlane Laboratories Limited) or monoclonal antibodies (Becton-Dickinson Transduction Laboratories), and anti-CREB or anti-phospho-CREB polyclonal antisera (1:500, Cell Signaling Technologies) overnight in 2% BSA, 0.01% Triton X-100, washed for 3 times ten minutes each in 2% BSA, 0.01% Triton X-100, stained for one hour at room temperature with goat-anti-rabbit Alexa Fluor 488, goat-anti-mouse Alexa Fluor 568, or goat-anti-mouse Alexa Fluor 594 (Molecular Probes), treated with DAPI for 5 minutes or To-Pro 3 (1:500, Molecular Probes) for 15 minutes, and then washed and mounted with gelvatol/DABCO (Sigma).

For phospho-CREB immunolocalization, cells or nuclei were fixed in 4% PFA for 30 minutes, permeabilized in 0.1% Triton X-100 for 15 minutes, blocked for at least one hour in 3% BSA, and immunostained using phospho-CREB polyclonal antisera (1:500, Cell Signaling Technologies) overnight at 4°C. Staining was visualized by incubation with goat-anti-rabbit Alexa Fluor 488 (Molecular Probes) for one hour at room temperature, treated with DAPI for 5 minutes, and then washed and mounted with gelvatol/DABCO (Sigma). Fluorescent images were recorded by a Hamamatsu digital camera connected to a Nikon inverted epifluorescent microscope. Images were taken at the same exposure time and gain, and all photographic manipulations were performed equally. Phospho-CREB positive nuclei were quantified in multiple fields from each stained slide by a blinded experimenter.

Confocal images were acquired with a Zeiss LSM 510 confocal system. Goat-anti-rabbit Alexa Fluor 488 was excited with a 488 nm Kr/Ar laser, goat-anti-mouse

Alexa Fluor 568 was excited with a 568 nm Kr/Ar Laser, and To-Pro 3 was excited with a 633 nm Kr/Ar Laser.

Isolation of Nuclei

Nuclei were isolated by cellular lysis followed by differential centrifugation (Spector et al., 1998) through OptiPrep (Axis-Shield). HeLa cells grown in suspension were lysed by detergent treatment in TM-2 buffer (0.01 M Tris-HCl, pH 7.4, 1.5 mM MgCl₂, 150 mM NaCl, 0.5 mM PMSF, 10 µg/ml Apoprotin, 10 µg/ml Leupeptin) containing 100 µg/ml digitonin followed by a 1000 x g spin. Supernatant (S1) was removed and the pellet was then resuspended in 0.25 M sucrose, 25 mM KCl, 30 mM MgCl₂, 20 mM Tris-HCl, pH 7.8. The resuspended pellet and 60% OptiPrep iodixanol were mixed (30% Optiprep final) and centrifuged at 10,000 x g for 20 minutes. The supernatant was removed and the nuclei-enriched pellet (P2) was resuspended in TM-2 buffer without detergent.

CREB Phosphorylation and Adenylyl Cyclase Assays

Equal aliquots of nuclei-enriched P2 preparations were incubated in 50 µl final volume of 100 mM Tris, pH 7.2, 10 mM MgCl₂, 5 mM ATP for CREB phosphorylation and 100 mM Tris, pH 7.2, 10 mM MgCl₂, 5 mM ATP, 0.5 mM IBMX for adenylyl cyclase assay with the indicated additions for 10 minutes (CREB phosphorylation) or 15 minutes (adenylyl cyclase) at 37° C. Reactions were stopped by addition of 20 µL of SDS sample buffer (CREB phosphorylation) or by being placed into a 100° C heat block for 3 minutes (adenylyl cyclase).

For whole cell and isolated nuclei CREB phosphorylation assays, equal cell or nuclear equivalents were separated under reducing conditions using a 10% SDS-PAGE, transferred to a PVDF membrane, and probed for CREB (rabbit polyclonal antiserum, Upstate Biotech) and phosphorylated CREB (rabbit polyclonal antiserum, Upstate Biotech). HRP conjugated secondary antibodies were used and bands were visualized using enhanced chemiluminescence. Fluorchem 8800 image analysis software (Alpha Innotech) was used to quantitate Western results. Intensities of phospho-CREB bands were normalized to total CREB.

cAMP produced in the cyclase assays was detected using a competition based assay with ^3H -cAMP (Amersham Pharmacia) and compared to a cAMP standard curve for quantitation.

Inhibitor profiles were determined by adenylyl cyclase assay (Assay Designs, Inc.) using purified sAC protein (Litvin et al., 2003) in the presence of 10 mM NaHCO_3 , 0.5 mM CaCl_2 , 10 mM MgCl_2 , and 10 mM ATP, or a mixture of purified catalytic domains, C1 and C2, from Type VII transmembrane adenylyl cyclase (Yan and Tang, 2002) in the presence of 5 mM MgCl_2 , and 1 mM ATP as previously described.

Quantitation of isolated nuclei (P2 fraction) immunocytochemistry

Nuclei were treated with Mg^{2+} -ATP alone or in combination with bicarbonate, forskolin, or 8-Br-cAMP as indicated above for 10 minutes, spread on a chilled slide, stored at -20°C , and then immunostained using phospho-CREB specific antisera as described. Nuclei were also treated with DAPI to differentiate intact nuclei from membrane ghosts. DAPI positive nuclei were scored for phospho-CREB immunofluorescence. Nuclei with detectable staining (i.e., NaHCO_3 panel, Figure 7A)

were considered positive for CREB phosphorylation whereas nuclei with no detectable staining (i.e., basal panel, Figure 7A) were counted as negative. Multiple microscopic fields were photographed for each condition, and data was combined from three to five separate experiments.

Western Analysis

Equal cell equivalents, unless otherwise noted, were separated under reducing conditions using a 10% SDS-PAGE, transferred to PVDF membrane, and blocked in 5% Milk. The blots were then probed with antibodies against either NaK ATPase (monoclonal, 1:50, Santa Cruz), histone H1 (monoclonal, 1:100, Santa Cruz), cytochrome oxidase subunit III (monoclonal, 2 μ g/ml, Molecular Probes), β -tubulin (monoclonal, 1:1000, Sigma), sAC (R21 monoclonal antibody, 1:500), monoclonal RI α or RII α antibodies (1:250, Becton Dickinson), or polyclonal RI α or RII α antisera (1:5000, Chemicon) overnight. HRP conjugated secondary antibodies were used and bands were visualized using enhanced chemiluminescence.

Acknowledgements

We thank Dr. Abijit Bapat for assistance quantifying nuclear phosphorylation of CREB, Kelley Rosborough for cryosectioning, Dr. Sylvia Finnemann for assistance with confocal microscopy, Dr. Yanqiu Chen for helpful suggestions, and Dr. Carl Nathan for critical reading of the manuscript. This work was supported by grants from the National Institutes of Health (AR32147 to D.A.F., GM62328 to J.B., HD42060 to J.B., HD38722 to L.R.L. and MSTP-GM07739 to J.H.Z.), Ellison Medical Foundation (to J.B.), American Diabetes Association (to J.H.Z.), and The Barbara and Stephen Friedman Fellowship Endowment (to J.H.Z.).

Abbreviations used in the paper:

AKAP, A kinase anchoring protein; cNGC, cyclic nucleotide gated channels; CRE, cAMP Response Element; CREB, CRE Binding Protein; DIC, differential interference contrast microscopy; EPAC, Exchange Protein Activated by cAMP; GPCR, G-protein coupled receptor; PDE, phosphodiesterase; PFA, paraformaldehyde; PKA, protein kinase A; sAC, soluble adenylyl cyclase; sAC_t, 48 kD isoform of sAC; tmAC, transmembrane adenylyl cyclase; Fsk, forskolin.

References

- Alto, N., J.J. Carlisle Michel, K.L. Dodge, L.K. Langeberg, and J.D. Scott. 2002. Intracellular targeting of protein kinases and phosphatases. *Diabetes*. 51 Suppl 3:S385-8.
- Bacskai, B.J., B. Hochner, M. Mahaut-Smith, S.R. Adams, B.K. Kaang, E.R. Kandel, and R.Y. Tsien. 1993. Spatially resolved dynamics of cAMP and protein kinase A subunits in Aplysia sensory neurons. *Science*. 260:222-6.
- Bailey, C.H., D. Bartsch, and E.R. Kandel. 1996. Toward a molecular definition of long-term memory storage. *Proc Natl Acad Sci U S A*. 93:13445-52.
- Bevensee, M.O., S.L. Alper, P.S. Aronson, and W.F. Boron. 2000. Control of intracellular pH. In *The Kidney*. Vol. I. D.W. Seldin and G. Giebisch, editors. Lippincott Williams & Wilkins, Philadelphia, PA.
- Buck, J., M.L. Sinclair, L. Schapal, M.J. Cann, and L.R. Levin. 1999. Cytosolic adenylyl cyclase defines a unique signaling molecule in mammals. *Proc Natl Acad Sci U S A*. 96:79-84.
- Byus, C.V., and W.H. Fletcher. 1982. Direct cytochemical localization of catalytic subunits dissociated from cAMP-dependent protein kinase in Reuber H-35 hepatoma cells. II. Temporal and spatial kinetics. *J Cell Biol*. 93:727-34.
- Chen, Y., M.J. Cann, T.N. Litvin, V. Iourgenko, M.L. Sinclair, L.R. Levin, and J. Buck. 2000. Soluble adenylyl cyclase as an evolutionarily conserved bicarbonate sensor. *Science*. 289:625-8.
- Constantinescu, A., I. Diamond, and A.S. Gordon. 1999. Ethanol-induced translocation of cAMP-dependent protein kinase to the nucleus. Mechanism and functional consequences. *J Biol Chem*. 274:26985-91.
- Daniel, P.B., W.H. Walker, and J.F. Habener. 1998. Cyclic AMP signaling and gene regulation. *Annu Rev Nutr*. 18:353-83.
- Davare, M.A., V. Avdonin, D.D. Hall, E.M. Peden, A. Burette, R.J. Weinberg, M.C. Horne, T. Hoshi, and J.W. Hell. 2001. A beta2 adrenergic receptor signaling complex assembled with the Ca²⁺ channel Cav1.2. *Science*. 293:98-101.
- De Cesare, D., and P. Sassone-Corsi. 2000. Transcriptional regulation by cyclic AMP-responsive factors. *Prog Nucleic Acid Res Mol Biol*. 64:343-69.
- Griffioen, G., P. Anghileri, E. Imre, M.D. Baroni, and H. Ruis. 2000. Nutritional control of nucleocytoplasmic localization of cAMP-dependent protein kinase catalytic and regulatory subunits in *Saccharomyces cerevisiae*. *J Biol Chem*. 275:1449-56.
- Hagiwara, M., A. Alberts, P. Brindle, J. Meinkoth, J. Feramisco, T. Deng, M. Karin, S. Shenolikar, and M. Montminy. 1992. Transcriptional attenuation following cAMP induction requires PP-1-mediated dephosphorylation of CREB. *Cell*. 70:105-13.
- Hagiwara, M., P. Brindle, A. Harootunian, R. Armstrong, J. Rivier, W. Vale, R. Tsien, and M.R. Montminy. 1993. Coupling of hormonal stimulation and transcription via the cyclic AMP-responsive factor CREB is rate limited by nuclear entry of protein kinase A. *Mol Cell Biol*. 13:4852-9.
- Hanoune, J., and N. Defer. 2001. Regulation and role of adenylyl cyclase isoforms. *Annu Rev Pharmacol Toxicol*. 41:145-74.
- Jaiswal, B.S., and M. Conti. 2003. Calcium regulation of the soluble adenylyl cyclase expressed in mammalian spermatozoa. *Proc Natl Acad Sci U S A*. 100:10676-81.
- Jungmann, R.A., M.R. Kuettel, S.P. Squinto, and J. Kwast-Welfeld. 1988. Using immunocolloidal gold electron microscopy to investigate cAMP-dependent protein kinase cellular compartmentalization. *Methods Enzymol*. 159:225-35.
- Kuettel, M.R., S.P. Squinto, J. Kwast-Welfeld, G. Schwoch, J.S. Schweppe, and R.A. Jungmann. 1985. Localization of nuclear subunits of cyclic AMP-dependent protein kinase by the immunocolloidal gold method. *J Cell Biol*. 101:965-75.
- Litvin, T.N., M. Kamenetsky, A. Zarifyan, J. Buck, and L.R. Levin. 2003. Kinetic properties of "soluble" adenylyl cyclase. Synergism between calcium and bicarbonate. *J Biol Chem*. 278:15922-6.
- Michel, J.J., and J.D. Scott. 2002. AKAP mediated signal transduction. *Annu Rev Pharmacol Toxicol*. 42:235-57.

- Murray, S.A., C.V. Byus, and W.H. Fletcher. 1985. Intracellular kinetics of free catalytic units dissociated from adenosine 3',5'-monophosphate-dependent protein kinase in adrenocortical tumor cells (Y-1). *Endocrinology*. 116:364-74.
- Pastor-Soler, N., V. Beaulieu, T.N. Litvin, N. Da Silva, Y. Chen, D. Brown, J. Buck, L.R. Levin, and S. Breton. 2003. Bicarbonate-regulated adenylyl cyclase (sAC) is a sensor that regulates pH-dependent V-ATPase recycling. *J Biol Chem*. 278:49523-9.
- Qiao, J., F.C. Mei, V.L. Popov, L.A. Vergara, and X. Cheng. 2002. Cell cycle dependent subcellular localization of exchange factor directly activated by cAMP. *J Biol Chem*. 277:26581-6.
- Riabowol, K.T., J.S. Fink, M.Z. Gilman, D.A. Walsh, R.H. Goodman, and J.R. Feramisco. 1988a. The catalytic subunit of cAMP-dependent protein kinase induces expression of genes containing cAMP-responsive enhancer elements. *Nature*. 336:83-6.
- Riabowol, K.T., M.Z. Gilman, and J.R. Feramisco. 1988b. Microinjection of the catalytic subunit of cAMP-dependent protein kinase induces expression of the c-fos gene. *Cold Spring Harb Symp Quant Biol*. 53:85-90.
- Rich, T.C., K.A. Fagan, H. Nakata, J. Schaack, D.M. Cooper, and J.W. Karpen. 2000. Cyclic nucleotide-gated channels colocalize with adenylyl cyclase in regions of restricted cAMP diffusion. *J Gen Physiol*. 116:147-61.
- Rich, T.C., K.A. Fagan, T.E. Tse, J. Schaack, D.M. Cooper, and J.W. Karpen. 2001. A uniform extracellular stimulus triggers distinct cAMP signals in different compartments of a simple cell. *Proc Natl Acad Sci U S A*. 98:13049-54.
- Sinclair, M.L., X.Y. Wang, M. Mattia, M. Conti, J. Buck, D.J. Wolgemuth, and L.R. Levin. 2000. Specific expression of soluble adenylyl cyclase in male germ cells. *Mol Reprod Dev*. 56:6-11.
- Singh, L.P., J. Andy, V. Anyamale, K. Greene, M. Alexander, and E.D. Crook. 2001. Hexosamine-induced fibronectin protein synthesis in mesangial cells is associated with increases in cAMP responsive element binding (CREB) phosphorylation and nuclear CREB: the involvement of protein kinases A and C. *Diabetes*. 50:2355-62.
- Spector, D.L., R.D. Goldman, and L.A. Leinwand. 1998. Culture and Biochemical Analysis of Cells. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Trumper, A., K. Trumper, and D. Horsch. 2002. Mechanisms of mitogenic and anti-apoptotic signaling by glucose-dependent insulinotropic polypeptide in beta(INS-1)-cells. *J Endocrinol*. 174:233-46.
- Wuttke, M.S., J. Buck, and L.R. Levin. 2001. Bicarbonate-regulated soluble adenylyl cyclase. *Journal of the Pancreas*. 2:154-8.
- Yan, S.Z., and W.J. Tang. 2002. Construction of soluble adenylyl cyclase from human membrane-bound type 7 adenylyl cyclase. *Methods Enzymol*. 345:231-41.
- Yang, J., J.A. Drazba, D.G. Ferguson, and M. Bond. 1998. A-kinase anchoring protein 100 (AKAP100) is localized in multiple subcellular compartments in the adult rat heart. *J Cell Biol*. 142:511-22.
- Zaccolo, M., and T. Pozzan. 2002. Discrete microdomains with high concentration of cAMP in stimulated rat neonatal cardiac myocytes. *Science*. 295:1711-5.
- Zhang, Q., D.W. Carr, K.M. Lerea, J.D. Scott, and S.A. Newman. 1996. Nuclear localization of type II cAMP-dependent protein kinase during limb cartilage differentiation is associated with a novel developmentally regulated A-kinase anchoring protein. *Dev Biol*. 176:51-61.
- Zippin, J.H., Y. Chen, P. Nahirney, M. Kamenetsky, M.S. Wuttke, D.A. Fischman, L.R. Levin, and J. Buck. 2003. Compartmentalization of bicarbonate-sensitive adenylyl cyclase in distinct signaling microdomains. *Faseb J*. 17:82-84.
- Zippin, J.H., L.R. Levin, and J. Buck. 2001. CO(2)/HCO(3)(-)-responsive soluble adenylyl cyclase as a putative metabolic sensor. *Trends Endocrinol Metab*. 12:366-70.

Figure Legends

Figure 1. Bicarbonate induces CREB phosphorylation via sAC activation in a PKA dependent manner. (A) COS7 cells were starved for bicarbonate 60 minutes, and were either incubated in the same bicarbonate starvation media for an additional 30 minutes (no treatment); incubated in normal, bicarbonate-containing DMEM in 5% CO₂ for 30 minutes (NaHCO₃); or preincubated with H89 (10 μM) for 10 minutes followed by incubation in normal, bicarbonate-containing DMEM in 5% CO₂ for 30 minutes (NaHCO₃ + H89). As control, COS7 cells grown in normal DMEM in 5% CO₂ were incubated with 10 μM forskolin for 30 minutes (forskolin). Cells were then immunostained with phospho-CREB antisera (green). Phase images of cells on left. Bar = 50 μm. (B) COS7 cells were transfected with vector control or sAC₁, and cells were assayed 36 hours post transfection. Cells were treated with vehicle control (DMSO) or H89 (10 μM) for 10 minutes and then stimulated with forskolin (10 μM) or given vehicle control (DMSO) for an additional 30 minutes. (C) Transfected cells were starved for bicarbonate, and were either incubated in the same bicarbonate starvation media for an additional 30 minutes; incubated in normal, bicarbonate-containing DMEM in 5% CO₂ for 30 minutes; or preincubated with H89 (10 μM) for 10 minutes followed by incubation in normal, bicarbonate-containing DMEM in 5% CO₂ for 30 minutes. (B & C) Top panel is Western blot using anti-phospho-CREB antisera; middle panel is Western blot using CREB-specific antisera with Phosphorylated CREB (P-CREB) protein and total CREB indicated. Shown below are the intensities of phospho-CREB relative to CREB normalized to vector control no treatment (lane 1).

Figure 2. Time course of CREB phosphorylation by sAC and tmAC. (A) Huh7 cells were starved for 1 hour for bicarbonate and CO₂ and then incubated in normal, bicarbonate-containing (44 mM) DMEM in 5% CO₂ for the time indicated or (B) kept in normal media and treated with PGE₂ (1 μM) for the time indicated. (A & B) Top panel is Western blot using anti-phospho-CREB antisera; middle panel is Western blot using CREB-specific antisera with Phosphorylated CREB (P-CREB) protein and total CREB indicated. Shown below are graphical representations of the intensities of phospho-CREB relative to CREB normalized to the 0 minute time point (lane 1).

Figure 3. Immunocytochemistry detects both sAC and PKA in the mammalian cell nucleus. (A) Confocal immunocytochemistry of Huh7 cells with monoclonal RIIα (center panels, green) and polyclonal RIα antibody (right panels, red). To-Pro 3 (upper left). Overlay of To-Pro 3 with both RIIα and RIα (lower left), RIIα (lower middle), and RIα (lower right). Secondary controls were negative (data not shown). (B-C) Confocal images of suspension HeLa cells immunostained with (B) PKA regulatory subunit RIIα polyclonal antisera and (C) PKA regulatory subunit RIα monoclonal antibody. Arrow A and B indicate suspension HeLa cytoplasm. Secondary controls were negative (insets). (D) Confocal immunocytochemistry of Huh7 cells stained with R41 monoclonal antibody against sAC.

Figure 4. Activated CREB , sAC, and PKA are present within the same rat liver nuclei. (A) Rat liver section stained with DAPI (DNA, blue, upper left), R52 biotinylated monoclonal antibody (sAC, green, upper middle), and polyclonal RIIα

antisera (R11 α , red, upper right), overlays of R11 α and sAC (lower left), sAC and DAPI (lower middle), and R11 α and DAPI (lower right). Arrows marked A indicate nuclei enriched for both sAC and PKA, whereas arrows marked B refer to nuclei not enriched for either. (B) Rat liver section stained with DAPI (DNA, blue, upper left), R21 monoclonal antibody (sAC, green, upper middle), and polyclonal P-CREB antisera (P-CREB, red, upper right), overlays of P-CREB and sAC (lower left), sAC and DAPI (lower middle), and P-CREB and DAPI (lower right). Arrows marked A indicate nuclei enriched for both sAC and P-CREB, whereas arrows marked B refer to nuclei enriched for neither. Rat liver tissue immunolocalization was confirmed to be inside the nucleus by confocal microscopy (data not shown).

Figure 5. sAC, PKA, and CREB coexist in mammalian cell nuclei. (A) Western blots of cell equivalents from HeLa whole cells (WC), low speed supernatant (S1), and nuclear-enriched high speed pellet (P2) probed with antibodies against NaK ATPase (NaK), histone H1 (Histone), cytochrome oxidase subunit III (COX), and β -tubulin (Tubulin). (B) Immunocytochemistry of nuclei isolated from HeLa cells (P2 pellet) using CREB polyclonal antisera (red) and sAC R52 biotinylated monoclonal antibody (green). DIC and DAPI (blue) images shown. Bar = 10 μ m. (C) Nuclei isolated from HeLa cells (P2 pellet) immunostained with polyclonal antisera (green) and monoclonal antibodies (red) directed against both R1 α and R11 α indicated that both proteins maintained their nucleoplasmic architecture throughout the fractionation procedure. Bottom row represents staining with goat-anti-rabbit (middle panel) or goat-anti-mouse controls (right panel) alone. Left column represents DAPI images in blue. Bar = 10 μ m. (D) Western blot of nuclear enriched P2 for sAC with R21 monoclonal antibody. (E)





Western blots of nuclear enriched P2 pellet with monoclonal (mRI α) and polyclonal (pRI α) antisera against RI α and with monoclonal (mRII α) and polyclonal (pRII α) antisera against RII α . All Westerns blots resolved only single bands of the predicted molecular weights.



Figure 6. sAC activity is the only detectable adenylyl cyclase activity in the mammalian cell nucleus.






(A) Adenylyl cyclase assay of whole cell lysate with Mg²⁺-ATP alone (Basal) or Mg²⁺-ATP and forskolin (Fsk). (B) Cyclase assay of nuclear lysate with Mg²⁺-ATP alone (Basal), or in the presence of 10 μ M forskolin (Fsk), or 40 mM bicarbonate (+NaHCO₃) in the absence (-) or presence of the sAC specific inhibitors KH1 (250 μ M) or KH2 (100 μ M). cAMP values are expressed as pmoles produced per ml of lysate and represent averages of duplicate determinations with standard deviation about the means indicated. These data are representative of at least 3 independent experiments performed in duplicate. (C) Chemical structures of inhibitors KH1 and KH2. (D) Inhibition of purified sAC (circles) or purified tmAC type VII C1 + C2 (squares) by KH1 (open symbols) and KH2 (filled symbols). Values for purified tmAC type VII C1 + C2 represent basal activity which was significantly above background.






Figure 7. Isolated nuclei contain a bicarbonate-responsive cAMP signaling microdomain dependent on both sAC and PKA. (A-B) Equal aliquots of nuclei enriched P2 were incubated with 40 mM NaCl (Basal), 10 μ M forskolin (FSK), or 40 mM sodium bicarbonate (NaHCO₃) for 10 minutes, smeared on a chilled glass slide, placed at -20°C, and immunostained for CREB family member phosphorylation using







phospho-CREB-specific polyclonal antisera (green). (A) Intact nuclei were confirmed by DAPI staining (left panels). Lower right panel highlights representative nuclei considered positive for CREB phosphorylation for quantitation. (B) Three microscopic fields per condition were photographed and counted by a blinded scientist. Values graphed represent the percentage of positive nuclei normalized to Basal (control) averaged from five separate experiments. Ratios above each bar represent the total number of positive nuclei divided by the total number of nuclei counted for all five experiments. (C) Western blot using phospho-CREB specific antisera against equal aliquots of nuclei-enriched P2 treated with Mg^{2+} -ATP alone (Basal), or substrate in the presence of 1 mM 8-Br-cAMP (cAMP), 40 mM bicarbonate, or 40 mM bicarbonate in the presence of either 10 μ M H-89 or 1 mM 8-Br-RpcAMPs. Each band was quantitated and normalized to basal; the relative intensities are basal (1 unit), cAMP (30 units), bicarbonate alone (27 units), bicarbonate plus H-89 (13 units), and bicarbonate plus Rp-cAMPs (8 units). (D) Western blot using phospho-CREB specific antisera against equal aliquots of nuclei-enriched P2 treated with 10 μ M forskolin (FSK) or with 40mM bicarbonate alone or in the presence of either sAC specific inhibitor KH1 (250 μ M) or KH2 (100 μ M).


Nam	screen #	Structure	In vitro IC50				In vivo IC50		Assay results
			pure sAC	pure sol. AC7	sAC in extracts	tmAC in extracts	sAC (basal)	tmAC (FSK)	
KH1	0782-6838		23 µM	> 200 µM			>>	>>	inhibits nuclear sAC inhibits pCREB MAY activate at ≤1 µM
KH2	2425-4155		2 µM	> 400 µM	>>100	>>100	>>	>>	inhibits nuclear sAC inhibits pCREB
KH3	2150-0290		8 µM	> 500 µM	>>100	>>100	>>	>>	inhibits pCREB
KH4	2150-0028		6 µM	>> 27 µM	>>100	>>100	>>	>>	inhibits pCREB
KH5	1683-2649		>> 55 µM	>> 55 µM	10-50 µM	100 µM	100 µM	>>	inhibits nuclear sAC inhibits pCREB
KH6	K074-5853		20 µM	>> 12 µM					

KH7	1185-0408		1 μ M	>> 55 μ M	10 μ M	>>100	10 μ M	> 500 μ M	inhibits nuclear sAC inhibits pCREB
KH8	2237-1900		0.5 μ M	0.5 μ M					
KH9									
KH10									

Nam	scr en #	Structure	In vitro EC50				In vivo IC50		Assay results
			pure sAC	pure tmAC	sAC in extracts	tmAC in extracts	sAC (basal)	tmAC (FSK)	
KH5	1683-2649		> 1 mM	>> 55 µM	10-100 µM	≥ 100 µM	100 µM	>>	inhibits nuclear sAC inhibits pCREB
KH7	1185-0408		1 µM	>> 55 µM	< 10 µM	>> 100 µM	10 µM - 30 µM	> 500 µM	inhibits nuclear sAC inhibits pCREB Inhibits insulin secretion
KH7.01	1185-0325		>100						
KH7.02	1185-0453		≥100				30 µM	≥200 µM	
KH7.03	2311-0174		≥300						

KH7.04	2311-0176		≥50						
KH7.05	2311-0184		~30				30 μM	>200 μM	
KH7.06	2311-0186		~30						
KH7.07	2311-0195		10				30 μM	>200 μM	
KH7.08	2311-0196		≥200				70 μM		





KH7.09	2311-0203		≥200						
KH7.10	2423-4824		80						
KH7.11	2423-4830		65						
KH7.12	2423-5085		30						
KH7.13	2423-5096		75						
KH7.14	3253-0230		>1mM					~200 μM	





KH7.15	3253-0233		~1mM					>200 μ M		INERT on Insulin secretion
--------	-----------	---	------	--	--	--	--	--------------	--	----------------------------




WHAT IS CLAIMED:

1. A method of treating a disorder in a subject mediated by soluble adenylyl cyclase, said method comprising:

administering to the subject an agent that inhibits soluble adenylyl cyclase, said agent being selected from the group consisting of compounds having the following formulae:









under conditions effective to treat the disorder mediated by soluble adenylyl cyclase.


2. The method according to claim 1, wherein the compound has the formula:




3. The method according to claim 1, wherein the compound has the formula:




4. The method according to claim 1, wherein the compound has the formula:




5. The method according to claim 1, wherein the compound has the formula:




6. The method according to claim 1, wherein the compound has the formula:




7. The method according to claim 1, wherein the compound has the formula:




8. The method according to claim 1, wherein the compound has the formula:




9. The method according to claim 1, wherein the compound has the formula:




10. The method according to claim 1, wherein the compound has the formula:



11. The method according to claim 1, wherein the compound has the formula:



12. The method according to claim 1, wherein the compound has the formula:



13. The method according to claim 1, wherein the disorder is selected from the group consisting of learning or memory disorders, male fertility/sterility, glaucoma, metabolic acidosis/alkalosis, diabetes, metabolic disorders, breathing disorders, and insulin resistance.

14. The method according to claim 13, wherein the disorder is a learning or memory disorder.



15. The method according to claim 13, wherein the disorder is male fertility/sterility.





16. The method according to claim 13, wherein the disorder is glaucoma.





17. The method according to claim 13, wherein the disorder is metabolic acidosis/alkalosis.


18. The method according to claim 13, wherein the disorder is diabetes.

19. The method according to claim 13, wherein the disorder is a metabolic disorder.
20. The method according to claim 13, wherein the disorder is a breathing disorder.
21. The method according to claim 13, wherein the disorder is an insulin resistance.
22. A method of inhibiting soluble adenylyl cyclase, said method comprising:
contacting eukaryotic cells with an agent that inhibits soluble adenylyl cyclase, said agent being selected from the group consisting of compounds having the following formulae:











under conditions effective to inhibit soluble adenylyl cyclase.


23. The method according to claim 22, wherein the compound has the formula:




24. The method according to claim 22, wherein the compound has the formula:




25. The method according to claim 22, wherein the compound has the formula:




26. The method according to claim 22, wherein the compound has the formula:




27. The method according to claim 22, wherein the compound has the formula:




28. The method according to claim 22, wherein the compound has the formula:




29. The method according to claim 22, wherein the compound has the formula:




30. The method according to claim 22, wherein the compound has the formula:




31. The method according to claim 22, wherein the compound has the formula:



32. The method according to claim 22, wherein the compound has the formula:



33. The method according to claim 22, wherein the compound has the formula:



Abstract

Bicarbonate responsive 'soluble' adenylyl cyclase resides, in part, inside the mammalian cell nucleus where it stimulates the activity of nuclear Protein Kinase A to phosphorylate the cAMP Response Element (CRE) Binding Protein. The existence of this complete and functional, nuclear-localized cAMP pathway establishes that cAMP signals in intracellular microdomains and identifies an alternate pathway leading to CRE Binding Protein activation.

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS



IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**


Zippin et al Figure 1




Zippin Figure 2




Zippin et al Figure 3




Zippin et al Figure 4




Zippin et al Figure 5



Zippin et al Figure 6



Zippin et al Figure 7



PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. EL984956509US

INVENTOR(S)					
Given Name (first and middle [if any])		Family Name or Surname		Residence (City and either State or Foreign Country)	
Jochen		Buck		4 Ferris Drive Greenwich, CT 06870	
Lonny R.		Levin		1161 York Avenue, Apt. 6D New York, NY 10021	
<input type="checkbox"/> Additional inventors are being named on the _____ separately numbered sheets attached hereto.					
TITLE OF THE INVENTION (280 characters max)					
CHEMICAL INHIBITORS OF SOLUBLE ADENYLYL CYCLASE (sAC)					
Direct all correspondence to: CORRESPONDENCE ADDRESS					
<input type="checkbox"/> Customer Number				Place Customer Number Bar Code Label here	
OR Type Customer Number here					
<input checked="" type="checkbox"/> Firm or Individual Name		Michael L. Goldman			
Address		Nixon Peabody LLP			
Address		Clinton Square, P.O. Box 31051			
City		Rochester	State	NY	ZIP 14603-1051
Country		USA	Telephone	(585) 263-1304	Fax (585) 263-1600
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification		Number of Pages		55	<input type="checkbox"/> CD(s), Number
<input type="checkbox"/> Drawing(s)		Number of Sheets			<input type="checkbox"/> Other (specify)
<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76					
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT					
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.				FILING FEE	
<input checked="" type="checkbox"/> A check or money order is enclosed to cover the filing fees				AMOUNT (\$)	
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge fees which may be required, or credit any overpayment to Deposit Account Number:				14-1138	
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.				\$80.00	
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.					
<input type="checkbox"/> No.					
<input checked="" type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: National Institutes of Health Grant Nos. HD38722, GM62328, and HD42060.					

Respectfully submitted,

Date 1/21/09

SIGNATURE

REGISTRATION NO.
(if appropriate)

30,727

TYPED or PRINTED NAME Michael L. Goldman

Docket Number:

19603/4650
(CRF D-3373-01)

TELEPHONE (585) 263-1304

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

Mail Stop Provisional Patent Application
Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450The PTO did not receive the following
listed item(s) 20 sheets of
claim

FEE TRANSMITTAL FOR FY 2003

Patent fees are subject to annual revision.

☒ Applicant claims small entity status. See 37 CFR 1.27

TOTAL AMOUNT OF PAYMENT (\$ 80

Complete if Known

Application Number	To Be Assigned
Filing Date	Herewith
First Named Inventor	Buck et al.
Examiner Name	
Art Unit	
Attorney Docket No.	19603/4650 (CRF D-3373-01)

METHOD OF PAYMENT (check all that apply)

☒ Check ☐ Credit Card ☐ Money Order ☐ Other ☐ None

☐ Deposit Account:

Deposit Account Number 14-1138

Deposit Account Name Nixon Peabody LLP

The Commissioner is authorized to: (check all that apply)

☐ Charge fee(s) indicated below ☒ Credit any overpayments

☒ Charge any additional fee(s)

☐ Charge fee(s) indicated below, except for the filing fee to the above-identified deposit account.

FEE CALCULATION

1. BASIC FILING FEE

Large Entity Fee Code	Large Entity Fee (\$)	Small Entity Fee Code	Small Entity Fee (\$)	Fee Description	Fee Paid
1001	770	2001	385	Utility filing fee	
1002	340	2002	170	Design filing fee	
1003	530	2003	265	Plant filing fee	
1004	770	2004	385	Reissue filing fee	
1005	160	2005	80	Provisional filing fee	80

SUBTOTAL (1) (\$ 80

2. EXTRA CLAIM FEES FOR UTILITY AND REISSUE

Total Claims	Extra Claims	Fee from below	Fee Paid
-20** =	X		0
Independent Claims -3** =	X		0
Multiple Dependent	X		0

Large Entity Fee Code	Large Entity Fee (\$)	Small Entity Fee Code	Small Entity Fee (\$)	Fee Description
1202	18	2202	9	Claims in excess of 20
1201	86	2201	43	Independent claims in excess of 3
1203	290	2203	145	Multiple dependent claim, if not paid
1204	86	2204	43	** Reissue independent claims over original patent
1205	18	2205	9	** Reissue claims in excess of 20 and over original patent

SUBTOTAL (2) (\$ 0

**or number previously paid, if greater; For Reissues, see above

FEE CALCULATION (continued)

3. ADDITIONAL FEES

Large Entity Fee Code	Large Entity Fee (\$)	Small Entity Fee Code	Small Entity Fee (\$)	Fee Description
1051	130	2051	65	Surcharge - late filing fee or oath
1052	50	2052	25	Surcharge - late provisional filing fee or cover sheet
1053	130	1053	130	Non-English specification
1812	2,520	1812	2,520	For filing a request for <i>ex parte</i> reexamination
1804	920*	1804	920*	Requesting publication of SIR prior to Examiner action
1805	1,840*	1805	1,840*	Requesting publication of SIR after Examiner action
1251	110	2251	55	Extension for reply within first month
1252	420	2252	210	Extension for reply within second month
1253	950	2253	475	Extension for reply within third month
1254	1,480	2254	740	Extension for reply within fourth month
1255	2,010	2255	1,005	Extension for reply within fifth month
1401	330	2401	165	Notice of Appeal
1402	330	2402	165	Filing a brief in support of an appeal
1403	290	2403	145	Request for oral hearing
1451	1,510	1451	1,510	Petition to institute a public use proceeding
1452	110	2452	55	Petition to revive - unavoidable
1453	1,330	2453	665	Petition to revive - unintentional
1501	1,330	2501	665	Utility issue fee (or reissue)
1502	480	2502	240	Design issue fee
1503	640	2503	320	Plant issue fee
1460	130	1460	130	Petitions to the Commissioner
1807	50	1807	50	Processing fee under 37 CFR 1.17(q)
1806	180	1806	180	Submission of Information Disclosure Stmt
8021	40	8021	40	Recording each patent assignment per property (times number of properties)
1809	770	2809	385	Filing a submission after final rejection (37 CFR 1.129(a))
1810	770	2810	385	For each additional invention to be examined (37 CFR 1.129(b))
1801	770	2801	385	Request for Continued Examination (RCE)
1802	900	1802	900	Request for expedited examination of a design application

Other fee (specify)

*Reduced by Basic Filing Fee Paid

SUBTOTAL (3) (\$ 0

CERTIFICATE OF MAILING OR TRANSMISSION [37 CFR 1.8(a)]

I hereby certify that this correspondence is being:

- ☐ deposited with the United States Postal Service on the date shown below with sufficient postage as first class mail in an envelope addressed to: Mail Stop _____ Commissioner for Patents, P. O. Box 1450, Alexandria, VA 22313-1450
- ☐ transmitted by facsimile on the date shown below to the United States Patent and Trademark Office at (703) _____

Date

Signature

Typed or printed name

SUBMITTED BY

Name (Print/Type) Michael L. Goldman

Signature

Registration No. (Attorney/Agent)

30,727

Complete (if applicable)

Telephone (585) 263-1304

Date

January 21, 2004

SEND TO: Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

EXPRESS MAIL CERTIFICATE

DOCKET NO.: 19603/4650 (CRF D-3373-01)
APPLICANTS: Jochen Buck and Lonny R. Levin
TITLE: CHEMICAL INHIBITORS OF SOLUBLE ADENYLYL
CYCLASE (sAC)

Certificate is attached to the **Provisional Application for Patent Cover Sheet (1 page) and Fee Transmittal (1 page)** of the above-identified application.

"EXPRESS MAIL" NUMBER: EL984956509US
DATE OF DEPOSIT: January 21, 2004

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to Mail Stop Provisional Patent Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

Shawn A. Lockett
(Typed or Printed Name of Person Mailing
Paper or Fee)


(Signature of Person Mailing Paper or Fee)